arXiv:1408.0995v2 [math.AG] 26 Nov 2014

A Genus Two Curve Related to the Class Number One Problem

by Viet Kh. Nguyen [1]

à la mémoire de V. A. Iskovskikh

Abstract

We give another solution to the class number one problem by showing that imaginary quadratic fields $\mathbb{Q}(\sqrt{-d})$ with class number $h(-d) = 1$ correspond to integral points on a genus two curve $\mathcal{K}_3$. In fact one can find all rational points on $\mathcal{K}_3$. The curve $\mathcal{K}_3$ arises naturally via certain coverings of curves: $\mathcal{K}_3 \to \mathcal{K}_6$, $\mathcal{K}_1 \to \mathcal{K}_2$ with $\mathcal{K}_2 \colon y^2 = 2x(x^3 - 1)$ denoting the Heegner curve, also in connection with the so-called Heegner-Stark covering $\mathcal{K}_1 \to \mathcal{K}_s$.

1. Introduction. In his famous *Disquisitiones Arithmeticae* (Article 303) Gauss hypothetically believed that there are only nine imaginary quadratic fields $\mathbb{Q}(\sqrt{-d})$ with class number $h(-d) = 1$, namely with $d = 1, 2, 3, 7, 11, 19, 43, 67, 163$. It is known nowadays as the Baker-Heegner-Stark theorem.

In the early 1980's V. A. Iskovskikh organized a seminar devoted to a beautiful approach of Heegner who was substantially first to give a complete proof of the theorem of Baker-Heegner-Stark based on an approach using modular functions ([2], *cf.* also [1], [6], [4], [7], [3], [5]). More precisely, assuming $d \equiv 3 \pmod 8$ (the main case), Heegner showed that every such a field corresponds to an integral point on the so-called Heegner curve:

$$(\mathcal{K}_2) \qquad y^2 = 2x(x^3 - 1)$$

In the higher class number case the statement reads as every field $\mathbb{Q}(\sqrt{-d})$, $d \equiv 3 \pmod 8$ corresponds to an integral point of degree $h(-d)$ on $\mathcal{K}_2$, provided $3 \nmid d$. One may have a remark that the "*exceptional*" field $\mathbb{Q}(\sqrt{-3})$ (corresponding to the point $(0, 0)$ on $\mathcal{K}_2$) is the only case with $3 \mid d$ such that j-invariant is a cube (namely equal to zero). On the contrary to the case of higher class numbers one sees, for example if $h(-d) = 2$, three fields $\mathbb{Q}(\sqrt{-51})$, $\mathbb{Q}(\sqrt{-123})$ and $\mathbb{Q}(\sqrt{-267})$ have no representatives on $\mathcal{K}_2$ along this line of arguments.

The aim of this note is to show that for our purpose one can involve another genus two curve $\mathcal{K}_3$:

$$(\mathcal{K}_3) \quad 8x^8 - 32x^6y + 40x^4y^2 + 64x^5 - 16x^2y^3 - 128x^3y + y^4 + 48xy^2 + 96x^2 - 32y - 24 = 0$$

[1] IMC Institute & Fpt University

2010 Mathematics Subject Classification. Primary 11G30, 11R29, 11R11.

realized at that time by the author.

The main result of the note is

Theorem A. *The curve $\mathcal{K}_3$ has the following integral points*

$$(3,6), \quad (-1,2), \quad (1,6), \quad (3,14), \quad (7,26), \quad (-17,150)$$

corresponding to the fields

$$\mathbb{Q}(\sqrt{-3}),\ \mathbb{Q}(\sqrt{-11}),\ \mathbb{Q}(\sqrt{-19}),\ \mathbb{Q}(\sqrt{-43}),\ \mathbb{Q}(\sqrt{-67}),\ \mathbb{Q}(\sqrt{-163})$$

respectively. Besides it has three "extraneous" integral points: $(-1,-2)$, $(-3,6)$, $(1,2)$ *(in fact* $(1,2)$ *is a double point on* $\mathcal{K}_3$*) plus two more rational points* $\big(-\frac{9}{17},\frac{6}{289}\big)$, $\big(-\frac{155}{79},\frac{42486}{6241}\big)$. *All the listed are the only rational points on* $\mathcal{K}_3$.

As a corollary one obtains a new solution to Gauss' "*tenth discriminant problem*".

Nowadays with the aid of computational tools one can easily compute the genus of $\mathcal{K}_3$. A surprising observation is the fact that $\mathcal{K}_3$ is birational to a genus 2 curve $\mathcal{K}_s$ studied independently by Heegner and Stark ([2], [6]).

Acknowledgments. I would like to thank V. A. Abrashkin for a nice talk on the class number problem at the seminar mentioned in the Introduction. This paper is dedicated to the 75th birthday of my teacher V. A. Iskovskikh with profound admiration and memory.

2. The curve $\mathcal{K}_3$. Let us first recall some known results from the Weber-Heegner-Birch theory ([10], [2], [1], [3]). We shall use Heegner's notation σ for the Schläfli functions of Stufe 48 which are defined by

$$\sigma(\tau) = q^{-\frac{1}{24}} \prod_{n=1}^{\infty} \big(1+q^{2n-1}\big); \quad \sigma_1(\tau) = q^{-\frac{1}{24}} \prod_{n=1}^{\infty} \big(1-q^{2n-1}\big);$$

$$\sigma_2(\tau) = \sqrt{2}\ q^{\frac{1}{12}} \prod_{n=1}^{\infty} \big(1+q^{2n}\big);$$

with $q = e^{i\pi\tau}$, $\mathrm{Im}(\tau) > 0$. Then the $24th$ powers σ^{24}, $-\sigma_1^{24}$, $-\sigma_2^{24}$ are the roots of

$$U^3 - 48U^2 + (768 - j)U - 4096 = 0 \tag{2.1}$$

where j is the modular invariant.

We assume throughout d is positive $\equiv 3 \pmod 8$ and put $\omega = \frac{1}{2}(3+\sqrt{-d})$. Then $-\sigma_2^{24}(\omega)$ is real and positive. Let ϕ denote the $3rd$ power $\sigma_2^3(\omega)$ up to a suitable $16th$ root of unity such that ϕ is real and positive. Then ϕ^8 is an algebraic integer satisfying equation (2.1) with $j = j(\omega)$ and by Weber's results ϕ^4, ϕ^2 are algebraic integers of degree $3h(-d)$. Furthermore Weber's conjecture proved in [1] tells us that indeed $\sqrt{2}\ \phi$ is integral of degree $3h(-d)$, so it satisfies a cubic equation

$$W^3 - 2a_3W^2 + 2b_3W - 8 = 0 \tag{2.2}$$

with a_3, b_3 integral of degree $h(-d)$. A computation in conjunction with (2.1) shows that (a_3, b_3) gives rise to an integral point of degree $h(-d)$ on $\mathcal{K}_3$. In particular if $h(-d) = 1$ we get the first statement of Theorem A.

If we argue in the same way with a cubic equation for ϕ^2

$$T^3 - 2a_2T^2 + 2b_2T - 8 = 0 \tag{2.3}$$

we get a relation between a_2, b_2 describing a curve $\mathcal{K}_6$ which is birationally isomorphic to the Pell conics

$$\Big(\frac{k}{2} - (a_2 + 1)\Big)^2 - 2\Big(\frac{a_2 + 1}{2}\Big)^2 = 1$$

where $b_2 = k(a_2 - 1) + 2$.

In view of (2.2), (2.3) the covering $\mathcal{K}_3 \to \mathcal{K}_6$ can be given in the following explicit form

$$a_2 = a_3^2 - b_3, \quad 2b_2 = b_3^2 - 8a_3 \tag{2.4}$$

from which may find interesting relations with the Pell sequence.

Remark 2.1. By virtue of (2.4) one sees that a_3 satisfies the following equation

$$z^4 - 2a_2z^2 - 8z + (a_2^2 - 2b_2) = 0 \tag{2.5}$$

Therefore (2.3) is nothing but Euler's resolvent for (2.5).

Example 2.2. As noted in the Introduction we illustrate our approach with some values d multiple of 3. By using the Tables from [10] and [9] one sees that the fields $\mathbb{Q}(\sqrt{-51})$, $\mathbb{Q}(\sqrt{-123})$, $\mathbb{Q}(\sqrt{-267})$ give rise to three real quadratic points

$$(-1, 8 + 2\sqrt{17}), \quad (4 + \sqrt{41}, 40 + 6\sqrt{41}), \quad (-10 - \sqrt{89}, 310 + 32\sqrt{89})$$

on $\mathcal{K}_3$ respectively.

3. The curve $\mathcal{K}_s$. If we assume in addition $3 \nmid d$, then the $8th$ powers σ^8, $-\sigma_1^8$, $-\sigma_2^8$ are the roots of

$$V^3 - \gamma_2 V - 16 = 0 \tag{3.1}$$

where γ_2 is the Weber function of Stufe 3 such that $\gamma_2^3 = j$.

In the notation of **2.** let φ be a cubic root of ϕ that is real and positive. Then as above φ^2, $\sqrt{2}\,\varphi$ are algebraic integral of degree $3h(-d)$. From (3.1) and a cubic equation for φ^2

$$Z^3 - 2\alpha_2 Z^2 + 2\beta_2 Z - 2 = 0 \tag{3.2}$$

with α_2, β_2 integral of degree $h(-d)$ we come to the equation $(\mathcal{K}_2)$ of Heegner's curve with $x = \alpha_2$, $y = \beta_2 - 2\alpha_2^2$.

Now writing a cubic equation for $\sqrt{2}\,\varphi$

$$S^3 - 2\alpha_3 S^2 + 2\beta_3 T - 4 = 0 \tag{3.3}$$

with α_3, β_3 integral of degree $h(-d)$ one gets a covering $\mathcal{K}_1 \to \mathcal{K}_2$ considered in [2], [6] which can be given by

$$\alpha_2 = \alpha_3^2 - \beta_3, \quad 2\beta_2 = \beta_3^2 - 4\alpha_3 \tag{3.4}$$

Hence the defining equation for $\mathcal{K}_1$ is as follows

$$(\mathcal{K}_1) \qquad 8x^8 - 32x^6y + 40x^4y^2 + 32x^5 - 16x^2y^3 - 64x^3y + y^4 + 24xy^2 + 24x^2 - 8y = 0$$

Despite a similarity with equation $(\mathcal{K}_3)$ this is a curve of genus 9.

Proposition 3.1. *The curve $\mathcal{K}_1$ has exactly 6 integral points $(0,0)$, $(1,2)$, $(-1,0)$, $(0,2)$, $(-1,2)$, $(2,6)$ which are in $1-1$ correspondence with class number one fields $\mathbb{Q}(\sqrt{-3})$, $\mathbb{Q}(\sqrt{-11})$, $\mathbb{Q}(\sqrt{-19})$, $\mathbb{Q}(\sqrt{-43})$, $\mathbb{Q}(\sqrt{-67})$, $\mathbb{Q}(\sqrt{-163})$.*

The proof follows immediately from (3.4) and [2]. The fact that these integral points exhaust all rational points on $\mathcal{K}_1$ follows from Heegner-Stark results ([2], [6]) on rational points on curve $\mathcal{K}_s$

$$(\mathcal{K}_s) \qquad w^2 = 2z(z^4 + 4z^3 - 2z^2 + 4z + 1)$$

The covering $\mathcal{K}_1 \to \mathcal{K}_s$ is given by

$$z = \frac{\beta_3}{\alpha_3^2} - 1; \quad w = 4(z-2)y - 2(3z^2 - 2z - 1), \quad y = \frac{1}{\alpha_3^3}$$

It should be noted that Heegner used a little different normalization (*cf.* [2], formulae $(151) - (152)$). Furthermore in order to enumerate all rational points of $\mathcal{K}_s$ one may apply the argument of [2] in solving equation $(\mathcal{K}_2)$ in integers. A precise statement is as follows.

Theorem 3.2. *The only rational points on the curve $\mathcal{K}_s$ are $(0,0)$, $(\pm 4, 1)$, $(\pm 4, -1)$, $\left(\pm \frac{7}{4}, \frac{1}{2}\right)$, $(\pm 14, 2)$.*

Theorem 3.3. *The curves $\mathcal{K}_3$ and $\mathcal{K}_s$ are birationally isomorphic.*

Since $\mathcal{K}_3$ is hyperelliptic of genus 2 one may use van Hoeij's algorithm ([8]) and a computing program (*e.g.* Maple) to write down its Weierstrass normal form which turns out to be the equation $(\mathcal{K}_s)$. The corresponding birational isomorphism $\mathcal{K}_s \to \mathcal{K}_3$ is given by

$$x = -\frac{z^4 + 8z^3 + 2wz + 18z^2 + 6w - 3}{z^4 + 4z^3 - 2z^2 - 12z + 1}$$

$$y = \frac{2\ P_8(z,w)}{(z^4 + 4z^3 - 2z^2 - 12z + 1)^2}$$

where

$$P_8(z,w) = (2z^5 + 10z^4 + 36z^3 + 68z^2 + 10z - 30)w$$
$$-z^8 + 60z^6 + 192z^5 + 82z^4 - 128z^3 + 172z^2 + 64z + 7.$$

Theorem A now follows from Theorems 3.2 and 3.3.

We note that the inverse birational isomorphism $\mathcal{K}_3 \to \mathcal{K}_s$ has more complicated formulae for (z, w). For completeness we include them here in the following form

$$z = 1 - \frac{4x^3 - 4xy - y^2 + 4x + 4}{2x^4 + 2x^3 - 3x^2y - 2xy + 6x - y + 2}$$

$$w = -\frac{2\ P_{12}(x, y)}{(x-1)(x^2+1)(x^2-2x-1)(x^2+2x+3)(x+1)^5}$$

where

$$\begin{aligned} P_{12}(x,y) \ = \ & 4x^{12} + 252x^{11} - 24x^{10}y + 156x^{10} - 622x^9y + 15x^8y^2 + 440x^9 - 514x^8y + \\ & +322x^7y^2 - x^6y^3 + 1256x^8 - 708x^7y + 288x^6y^2 - 21x^5y^3 + 1536x^7 - 620x^6y \\ & +310x^5y^2 - 19x^4y^3 + 1344x^6 - 716x^5y + 64x^4y^2 - 20x^3y^3 + 440x^5 - 640x^4y \\ & +22x^3y^2 - 7x^2y^3 - 12x^4 - 316x^3y - 8x^2y^2 - 3xy^3 - 124x^3 - 140x^2y - 6xy^2 \\ & -y^3 - 92x^2 - 22xy + y^2 - 16x + 2y. \end{aligned}$$

Remark 3.4. There is a map $\mathcal{K}_2 \to \mathcal{K}_6$ ([2]) given by

$$a_2 = 4\alpha_2^3 - 6\alpha_2\beta_2 + 3; \quad b_2 = 4\beta_2^3 - 12\alpha_2\beta_2 + 6$$

and similarly a map $\mathcal{K}_1 \to \mathcal{K}_3$

$$a_3 = 2\alpha_3^3 - 3\alpha_3\beta_3 + 3; \quad b_3 = \beta_3^3 - 6\alpha_3\beta_3 + 6$$

which turned out quite useful in the computation process.

References

[1] Birch, B. J., *Weber's class invariants*, Mathematika **16** (1969), 283–294.

[2] Heegner, K., *Diophantische Analysis und Modulfunktionen*, Math. Z. **56**, Heft 3 (1952), 227-253.

[3] Schertz, R., *Complex multiplication*, Cambridge University Press, 2010.

[4] Serre, J.-P., *Lectures on the Mordell-Weil theorem*, Vieweg, 1989.

[5] Shafarevich, I. R., *The tenth discriminant problem*, Algebra i Analiz, **25**:4 (2013), 260-277 (in Russian).

[6] Stark, H. M., *Class numbers of complex quadratic fields*, Modular Functions of One Variable I, Ed.: Kuyk W., Lect. Notes Math. **320** (1973), 153–174.

[7] Stark, H. M., *The Gauss class-number problems*, Clay Math. Proc., **7** (2007), 247-256.

[8] van Hoeij, M., *An algorithm for computing the Weierstrass normal form of hyperelliptic curves*, arXiv:math/0203130, www.math.fsu.edu/∼hoeij/algcurves.html

[9] Watson, G. N., *Singular moduli* (5), Proc. London Math. Soc., **42** (1937), 377-397.

[10] Weber, H., *Elliptische Functionen und algebraische Zahlen*, Braunschweig, 1891; and *Lehrbuch der Algebra*, 2-te Auflage, Bd 3, Braunschweig, 1908.

IMC Institute & Fpt University
Permanent: 8 Ton That Thuyet, My Dinh, Tu Liem, Hanoi, Vietnam
e-mail: vietnk@fpt.edu.vn, nhimsocvn@gmail.com